團聚健康檢查證明	檢查日期/// (月)(日)(年)	
附表 (醫院名稱、地址、電話、傳真機) (健檢醫院可另行開立具有醫院標誌之健康證明)		
基本資料 (BASIC DATA)		
姓名: 性別: □男 Male □女 Female Name Sex : □別 Male □□ 女 Female 身分證字號: : : : ID No. : : :	照片	
超 NO. 護照號碼 . Passport No. Date of Birth //	Photo	
實驗室檢查(LABORATORY EXAMINATIONS)		
 A. HIV 抗體檢查(Serological Test for HIV Antibody): □陽性(Positive) □陰性 □未確定(Indeterminate) a.篩檢(Screening Test): □EIA □Serodia □其他(Others) b.確認(Confirmatory Test): □Western Blot □其他(Others) B.胸部 X 光檢查肺結核(Chest X-Ray for Tuberculosis): (妊娠孕婦可免接受「胸 □上常(Normal) □異常(Abnormal)※限大片攝影(St C.腸內寄生蟲(含痢疾阿米巴等原蟲)糞便檢查(Stool examination for parasites ind <i>histolytica</i> etc.): □陽性, 種名(Positive, Species) D. 梅毒血清檢查(Serological Test for Syphilis): □陽性(Positive) □陰性(Neg a. □RPR b.□VDRL c.□TPHA/TPPA d.□其它(Other) E. 申請者應檢具麻疹、德國麻疹(風疹)抗體陽性檢驗報告或提供麻疹、德國麻疹預時 should provide proof of positive measles and rubella antibody titers or measles and rub certificates): a. □抗體檢查(Antibody test)麻疹抗體 measles antibody titers □陽性 Posit 德國麻疹預防接種證明 Vaccination Certificates of Measles □德國麻疹預防接種證明 Vaccination Certificates of Rubella c. □經醫師評估,有接種禁忌者,暫不適宜接種。(Having contraindications, not s 	m X 光檢查」) andard Film Only) cludes Entameba 陰性 (Negative) gative) 防接種證明(Applicant bella vaccination ive □陰性 Negative ive □陰性 Negative	
漢 生 病 檢 查(CHECK-UP FOR Hanser 漢生病(麻風病)視診結果(Skin Check-up) □正常 Normal □異常 Abnormal (※視診 確認) (※If abnormal skin lesion is found, further skin biopsy or skin smear is required a.病理切片(Skin Biopsy): □陽性(多菌、少菌性【Positive - MB,PB】; 診斷依據	異常者,須進一步採檢)	
【Diagnostic if either of them positive】)		
 備註: 一、本表供大陸地區人民首次申請來臺團聚時使用。This form is for use in applying 二、妊娠孕婦可免接受「胸部 X 光檢查」。A pregnant woman is not necessary to hav 三、梅毒血清檢查陽性者,檢具治療證明,視為合格。Those who have positive ress syphilis should submit medical treatment certificate. 四、健康檢查證明不合格之認定原則詳如附錄。Appendix is principles in determinit 五、根據以上對 先生/女士/小姐之檢查結果為□合格 □不合格。 Conclusion: This is to certify that, based on the above medical report, Mr/Mrs/Ms He/She □passes □fails the checkup. 	e chest X-ray examination sults for serological test for ing the health status failed.	
負責醫師簽章 (Chief Physician):		
醫院負責人簽章(Superintendent):		
日期 (Date):/ ※本證明三個月內有效 (Vaild for Three M	Ionths) 01/01/2009	

附錄:	健康檢查證明不合格之認定原則
-----	----------------

檢驗項目	不合格之認定原則
人類免疫缺乏病	一、人類免疫缺乏病毒抗體檢驗經初步測試,連續二次呈陽性反應者,應以西方墨點法(WB)作確
毒抗體檢查	認試驗。
	二、連續二次(採血時間需間隔三個月)西方墨點法結果皆為未確定者,視為合格。
胸部X光檢查	一、活動性肺結核(包括結核性肋膜炎)視為「不合格」。
	二、非活動性肺結核視為「合格」,包括下列診斷情形:纖維化(鈣化)肺結核、纖維化(鈣化)病灶及
	肋膜增厚。
腸內寄生蟲糞便	一、經顯微鏡檢查結果為腸道蠕蟲蟲卵或其他原蟲類如:痢疾阿米巴原蟲(Entamoeba
檢查	histolytica)、鞭毛原蟲類,纖毛原蟲類及孢子蟲類者為不合格。
	二、經顯微鏡檢查結果為人芽囊原蟲及阿米巴原蟲類,如:哈氏阿米巴 (Entamoeba hartmanni)、
	大腸阿米巴(Entamoeba coli)、微小阿米巴(Endolimax nana)、嗜碘阿米巴(Iodamoeba
	butschlii)、雙核阿米巴(Dientamoeba fragilis)等,可不予治療,視為「合格」。
梅毒	一、以 RPR 或 VDRL 其中一種加上 TPHA(TPPA)之檢驗,如檢驗結果有下列情形任一者,為「不
	合格」:
	(一)活性梅毒:同時符合通報條件(一)及(二)、或僅符合通報條件(三)者。
	(二)非活性梅毒:僅符合通報條件(二)者。
	二、通報條件:
	(一)臨床症狀出現硬下疳或全身性梅毒紅疹等臨床症狀。
	(二)未曾接受梅毒治療或病史不清楚者,RPR(+)或 VDRL(+),且 TPHA (TPPA)=1:320 以上(含
	320) •
	(三)曾經接受梅毒治療者,VDRL 價數上升四倍。
麻疹、德國麻疹	麻疹、德國麻疹(風疹)抗體陰性且未檢具麻疹、德國麻疹預防接種證明者為不合格。但經醫師評估
(風疹)	有麻疹、德國麻疹疫苗接種禁忌者,視為合格。

Appendix: Principles in determining the health status failed

Test Item	Principles on the determination of failed items
	1. If the preliminary testing of the serological test for HIV antibody is positive for two consecutive times,
for HIV	confirmation testing by WB is required.
Antibody	2. When findings of two consecutive WB testing (blood specimens collected at an interval of three
5	months) are indeterminate, this item is considered qualified.
Chest X-ray	1. Active pulmonary tuberculosis (including tuberculous pleurisy) is unqualified.
5	2. Non-active pulmonary tuberculosis including calcified pulmonary tuberculosis, calcified foci and
	enlargement of pleura, is considered qualified.
Stool	1. By microscope examination, cases are determined unqualified if intestinal helminthes eggs or other
Examination for	
Parasites	2. Blastocystis hominis and Amoeba protozoa such as Entamoeba hartmanni, Entaboeba coli, Endolimax
	nana, Iodamoeba butschlii, Dientamoeba fragilis found through microscope examination are
	considered qualified and no treatment is required.
Serological Test	1. After testing by either RPR or VDRL together with TPHA(TPPA), if cases meet one of the following
for Syphilis	situations are considered failing the examination.
	(1)Active syphilis: must fit reportable criterion $(1) + (2)$ or only reportable criterion (3) .
	(2)Inactive syphilis: only fit reportable criterion (2).
	2. Reportable criterion:
	(1)Clinical symptoms with genital ulcers (chancres) or syphilis rash all over the body.
	(2)No past diagnosis of syphilis, a reactive nontreponemal test (i.e., VDRL or RPR), and
	$TPHA(TPPA) = 1 : 320\uparrow(including 1 : 320)$
	(3)A past history of syphilis therapy and a current nontreponemal test titer demonstrating four fold or
	greater increase from the last nontreponemal test titer.
Measles,	The item is considered unqualified if measles or rubella antibody is negative and no measles, rubella
Rubella	vaccination certificate is provided. Those who having contraindications, not suitable for vaccinations are
	considered qualified.